



# POISON

KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

# TE 8 PLUS

## FERTILIZERS

### GUARANTEED ANALYSIS

Nitrogen (N)	2.6% w/v
N as nitrate	2.6% w/v
Potassium (K)	0.1% w/v
Sulphur (S)	4.2% w/v
Magnesium (Mg)	2.4% w/v
Manganese (Mn)	3.2% w/v
Zinc (Zn)	3.2% w/v
Copper (Cu)	0.5% w/v
Molybdenum (Mo)	0.02% w/v
Boron (B)	0.2% w/v
Iron (Fe)	0.7% w/v
Fulvic Acid	0.5% w/v
Specific Gravity @ 20°C	1.295 kg/L
pH	1.0 to 2.0

(Formerly TE 6 PLUS)  
Multitrace blend containing S, Mg, Mn, Zn,  
Cu, Fe, B, Mo & Fulvic Acid  
SNPK0046  
Zinc Sulphate 83 g/L

**WARNING:**  
KEEP OUT OF REACH OF CHILDREN



20 L



## APPLICATION RATES

### Typical Fertigation (via irrigation systems)

#### Application Rates:

Apply at 10 to 25 L/ha toward the end of the irrigation shift. Know your injection lag times and allow adequate time, following the injection of the product, to flush the irrigation system out with clean water, including all mainlines, submains and lateral delivery systems.

### Typical Application Rates

#### (At Planting, Directed Soil Spray):

20 to 30 L/ha as a directed soil spray, prior to planting or banded under canopy, with 100 to 400 L/ha water, or irrigated in after application (depending on crop).

### Typical Foliar Application Rates:

2 to 10 L/ha  
Horticulture use 200 to 2,000 L/ha water  
Broadacre use at least 100 L/ha water

## GUIDING PRINCIPLES

Before using TE 8 PLUS, seek appropriate agronomic advice via SLTEC's Balanced Agronomy services and dealer agronomist network.

Soil analysis combined with plant tissue (or SAP) analysis are recommended to assist in making an informed decision about which product best meets your crop, soil and general environmental conditions.

Fertilizer can burn and or damage crops and pasture. Because climatic and soil conditions, application methods, irrigation and management practices are beyond the control of Sustainable Liquid Technology Pty Ltd and cannot be foreseen, Sustainable Liquid Technology Pty Ltd accepts no responsibility whatsoever for any commercial damage, loss or other result following the use of this product, whether used in accordance with directions or not, subject to any overriding statutory provision and provided that such liability under those provisions shall be limited to the replacement of the goods as supplied or the rendering again of the services that are provided. The buyer accepts and uses this product subject to these conditions.

Batch No. or Tank Number & Date:

PO Box 43 / 2055 Finlay Road  
TONGALA VICTORIA 3621  
P: 1800 768 224 F: 03 5859 1363  
E: enquiries@sltec.com.au  
ABN: 63 234 073 378 ACN: 113 670 269  
[www.sltec.com.au](http://www.sltec.com.au)

## PRODUCT INFORMATION

### PRODUCT STORAGE

Keep material in a cool, dry, well ventilated area, away from direct sunlight, between 10 to 30°C. Store product in suitable containers made from material such as stainless steel, high density polyethylene and fibreglass. Fluid fertilizers can be corrosive to material such as brass, copper, zinc or alloys of these metals. Clean equipment after use thoroughly with clean fresh water. Use SLTEC's Fert Tank Cleaner to rinse fertilizer tanks and transfer systems. TE 8 PLUS is recommended for use within 6 months from dispatch date.

### WARNING & SAFETY INFORMATION

**WARNING:** Do not swallow. The spray from this product may act as an irritant. Avoid inhalation and contact with the eyes and skin. Wear rubber gloves and appropriate eye protection when handling the product. See the product MSDS for more information, [www.sltec.com.au](http://www.sltec.com.au)

**WARNING:** Stock should not be allowed to graze the pasture for at least three weeks following application, to avoid the potential risk of nitrate poisoning and to maximise the growth response to the applied nitrogen. Do not cut for silage or hay during this with holding period also.

**WARNING:** This product contains heavy metal impurities. Its use may result in accumulation of cadmium in the soil. Depending on soil characteristics, irrigation water quality, plant species and variety, crop uptake of cadmium may lead to residue levels in plant and animal products in excess of the maximum level specified by the Australia New Zealand Food Standards Code. In pasture, the offfal from grazing animals may also exceed these limits.

**First Aid:** For advice, contact a Poisons Information Centre (phone Australia 131 126) or a doctor. If swallowed, do NOT induce vomiting. If in eyes, hold eyelids apart and flush the eye continuously with running water. Continue flushing until advised to stop by a Poisons Information Centre (phone Australia 131 126) or a doctor, or for at least 15 min. If skin or hair contact occurs, remove contaminated clothing and flush skin and hair with running water.

**Safety Directions:** Avoid contact with eyes. Avoid contact with skin.

### IMPURITIES

Cadmium (Cd) 1.7 mg/kg

Dispatch Date:

